

## Sexual Dysfunction in Men with Systolic Heart Failure and Associated Factors

Sharareh Zeighami Mohamadi\*<sup>MSc</sup>, Mandana Shahparian<sup>1MD</sup>, Farzad Fahidy<sup>2 BSc</sup>, Ebrahim Fallah<sup>2 BSc</sup>

\*. *Instructor of Nursing, School of Nursing and Midwifery, Islamic Azad University of Karaj, Iran*

1. *Specialist of cardiovascular disease, social security Alborz hospital, Karaj, Iran*

2. *Nursing graduate, social security Shahriar hospital, Shahriar, Iran*

### Abstract

**Aims:** Heart failure is a major risk factor for erectile dysfunction. This study seeks to investigate the incidence and factors associated with erectile dysfunction in patients with systolic heart failure.

**Methods:** This is a cross sectional study that was conducted on 100 male patients with systolic heart failure in 2009. Convenience sampling was used to gather the required data. Data collection instruments included a demographic data form, IIEF-5 (The International Index of Erectile Function, a 5-item version), MLHFQ (Minnesota Living with Heart Failure), and CES-D (Center for Epidemiologic Studies Depression Scale). The data was analyzed by SPSS software and using descriptive statistics, t-test, ANOVA and Pearson correlation coefficient.

**Findings:** The findings indicated that the mean score of erectile dysfunction was  $14.02 \pm 6.26$  and 80% of male patients with systolic heart failure suffered from erectile dysfunction. There was a significant relationship between erectile dysfunction and age ( $p < 0.001$ ), education ( $p = 0.019$ ), occupation ( $p = 0.002$ ), hemoglobin level ( $p = 0.003$ ), left ventricular ejection fraction ( $p = 0.030$ ), cholesterol level ( $p = 0.001$ ), renal dysfunction ( $p = 0.009$ ), use of digoxin ( $p = 0.014$ ), angiotensin converting enzyme inhibitors ( $P < 0.001$ ), beta blocker ( $p = 0.001$ ), diuretics ( $p = 0.035$ ), depression ( $p < 0.001$ ), and quality of life ( $p < 0.001$ ).

**Conclusion:** Erectile dysfunction is the most common problem in patients with systolic heart failure and is associated with age, medical conditions, co-morbidity, medicinal treatments and psychological disorders. In patients with heart failure, erectile dysfunction has a negative impact on quality of life.

**Keywords:** Heart failure, Erectile dysfunction, Depression, Quality of life

### Introduction

Heart failure is a clinical syndrome resulted from heart inability to pump an essential amount of oxygenated blood which is created to satisfy metabolic needs of body. There are two types of systolic and diastolic heart failure. Systolic heart failure is caused by heart inability to pump enough blood while diastolic heart failure is a result of inability of ventricles to relax and get filled during diastole [1]. About five million people in the United States of America are suffering from heart failure and five hundred thousand new cases are added to them every year, too [2]. Heart inability to circulate blood creates several symptoms such as being short of breath, fatigue, dizziness, angina pectoris, edema, and ascites [3]. Intolerance of labors, getting exhausted, and being short of breath disturb patients' quality of life and functional ability [4]. They cause patients to be unable to fulfill their duties and therefore face financial problems [5], experience disturbance in their ordinary activities of life, lose their independence and depend on others to perform self-care activities, get isolated from

the society, face sexual dysfunction, and change their family and social roles in life [3]. Sexual function is an important aspect of quality of life. Sexual dysfunction affects the quality of life of patients with heart failure and reduces it [6], and results in a decrease in libido, reduction of number of intercourses, and dissatisfaction with sexual intercourse [7]. Heart failure itself is known as a risk factor for erectile dysfunction. Erectile dysfunction is the inability to achieve or maintain an erection for a satisfactory intercourse with sex partner [8]. In a survey conducted by Medina, 74% [9], Schwartz, 84% [10], and Rastogi, 75% of patients with heart failure had erectile dysfunction [11]. Erectile dysfunction is caused as a result of mental, neurological, hormonal, or vascular problems or side effects of drugs [7]. Various factors are involved in erectile dysfunction in men with heart failure such as ageing [12, 13], smoking [14], Body Mass Index (BMI) [15], suffering from other chronic diseases such as hypertension, diabetes, chronic obstructive pulmonary diseases, ischemic heart diseases, hyperlipidemia, anemia [16 and 17], intensity of heart failure and left

\* Correspondence Author: zeighami@kia.ac.ir

ventricular ejection fraction [18], and side effects of medications such as digoxin, beta-blocker, diuretics and spironolactone [11]. Healthy sexual activity plays an important role in men and women's mental health, family formation and stability, and preventing disagreements, misunderstandings, marital breakdown, and its unfortunate consequences [19]. In the past decade, social attitude has changed towards sexual health and hygiene [20]. Nowadays, many scholars and experts increasingly emphasize on identifying and removing people's sexual problems [19]. Considering rapid growth in the number of patients with chronic heart failure, it is essential to know the effective factors in sexual disorders which can damage quality of life. After identifying them, it is necessary to remove the problems via using different methods to increase the quality of life in those patients. This study was carried out to investigate the incidence of erectile dysfunction and factors associated with it in patients with systolic heart failure.

### Methods

This is a cross sectional study which investigated erectile dysfunction and associated factors with it. The research population included all patients with systolic heart failure hospitalized in the internal ward and clinic of Alborz Ta'mine Ejtema'i in Karaj and Ta'mine Ejtema'i Hospital in Shahriar. It was conducted from June 2010 to December 2010. Research samples included a hundred patients with heart failure which were chosen from the research population through convenience sampling method and regarding the selection criteria (married, having heart failure for at least one year, ejection fraction below 40 %, not having previous records of mental diseases, depression, or sterility). The researcher assistant visited the mentioned hospitals every day. Considering the selection criteria, after explaining the purpose of the study, he completed and collected the research instruments through interviews. The instruments used in this study included sample selection form, demographic

information form with some information about the disease (age, marital status, education, income level, family economic status, occupation, smoking, disease duration, type of chronic disease along with heart failure, type of used medications, left ventricular ejection fraction based on echocardiography, BMI, serum hemoglobin level and cholesterol), IIEF-5, MLHFQ, and CES-D. In the present study, smoking meant smoking at least one cigarette a day for one year. BMI was calculated through dividing weight in kilos by the square of height in meters. To get participants' height, they were asked to remove their shoes and stand next to a wall with their back on it while they closed their legs tight and their knees, pelvis, shoulder and head were along each other. Afterwards, the researcher put the tape measure next to participants' head and got their height by one centimeter accuracy.

IIEF-5 contained 5 questions. Its answers were graded from one to five scores according to 5-point likert scale. The score ranged from 5 to 25. A score below 21 suggested erection failure. Finally, based on the obtained scores, participants were classified into four categories: high erection failure (5 to 10), average erection failure (11 to 15), low erection failure (16 to 20), and normal condition (21 to 25). The reliability was found out to be 0.96 [21].

MLHFQ has been designed to obtain information about the quality of life of patients with heart failure. This specific questionnaire was designed by Rector in 1984 to identify the effect of treatments on the quality of life of patients with heart failure. It is the most common instrument in evaluating the quality of life in these patients in research studies. This questionnaire shows patients' understanding of the impacts of heart failure on physical, economical, social, and psychological aspects of life. Questions of this questionnaire deal with symptoms of the disease such as being short of breath, fatigue, peripheral edema, sleep disorders and psychological symptoms such as depression, anxiety, social relationships, physical and sexual activities, work, and emotions. Each

Table 1: Investigation of erectile dysfunction score correlation with the studied variables in men with systolic heart failure

Age	p<0.001	r=-0.417
Income	p=0.576	r=0.057
Disease duration	p=0.133	r=-0.151
Blood cholesterol level	P=0.001	r=-0.322
Hemoglobin	P=0.003	r=0.295
Left ventricular ejection fraction	P=0.030	r=0.217
BMI	P=0.234	r=0.120
Depression	p<0.001	r=-0.608
Quality of life	p<0.001	r=-0.556

question has six criteria which are graded from zero to five scores. Zero represents the best condition and number five shows the worst condition. Therefore, the higher the total score is, the lower the quality of life is. Validity and reliability of this questionnaire is higher than other questionnaires. In Rector's study, reliability of this instrument was reported to be 0.94 [22].

CES-D: This questionnaire contains 20 questions on depression. It is related to Center for Epidemiologic Studies and depression level is graded through 20 questions with 4-point likert scale from zero to three. In this study, 0 = rarely or never (less than a day), 1 = slightly (one to two days), 2 = Sometimes (three to four days), 3 = usually (five to seven days). Based on the total scores obtained from this instrument, a score of less than 15 means no depression, 15 to 21 equals little to mild depression, and a score above 21 suggests severe depression [23, 24]. Reliability of this questionnaire was 0.85 in Clark's study [25]. In order to determine the scientific reliability of the above mentioned instruments, content validity method was used through the study of scientific sources and according to the research objectives. To identify the reliability of the questionnaire in the present study, test-retest method was used. In this method,

questionnaires were given to 10 patients with heart failure in 2 phases in a 10-day interval and the correlation of the two tests was estimated. In this study, for IIEF-5  $r = 0.78$ , for MLHFQ  $r = 0.70$ , and for CES-D  $r = 0.84$ . Moral issues in this study were taken into consideration. The hospital authorities' permission for the study was taken, a formal letter of introduction was presented to the hospital authorities, the researcher was introduced to the participants and the purpose and method of the study were explained to them, all the participants gave written consent for attending the study and having the right to leave it, it was attempted that the study did not violate the culture, traditions, and customs of the society, and the authorities and patients were ensured that the results of the study would be announced if they liked.

To analyze the data, SPSS software version 14 was utilized. The data was described by frequency tables, mean and SD. For data analysis, Pearson correlation coefficient, t-test for independent groups and ANOVA were used. Significance level in this study was considered to be 0.05.

## Findings

The mean of age of men with systolic heart failure was  $59.23 \pm 11.69$  with a range of 34 to 88 years old, and the mean of duration of heart failure was  $4.59 \pm 2.95$  years with a range of 1 to 15 years. The mean of monthly income was  $486000 \pm 254.729$  tomans with a range of 200000 to 2400000 tomans per month. 46% of participants had an average economic condition, 42% of them had primary education, and 51% were retired. Their mean of BMI was  $26.18 \pm 4.47$  with a range of 18.44 to 38.52, the mean of left ventricular ejection fraction was  $35.19 \pm 7.43$  with a range of 10 to 40%, the mean of hemoglobin level was  $14.32 \pm 1.86$  with a range of 9.9 to 20.3 g/dl, and the mean of blood cholesterol level was  $171.98 \pm 43.21$  with a range of 103 to 327. 62% of participants never smoked. The most common co-morbidity was hypertension (33%) and diabetes (33%). 41% of participants were being treated with digoxin, 38% of them with diuretics, 29% with

angiotensin converting enzyme inhibitors, 5% with calcium blocker, 19% with beta blockers, 74% with nitrate, 35% with angiotensin blockers, and 8% with aldactone. The mean of erectile dysfunction score was  $14.02 \pm 6.26$  and 80% of men with systolic heart failure had low intensity erectile dysfunction. The mean of depression score in men with heart failure was  $20.31 \pm 12.06$ , and the mean of total score of quality of life was  $41.61 \pm 21.30$ .

There was a significant difference between the mean of erectile dysfunction score and education ( $p=0.019$ ) and job ( $p=0.002$ ); however, no meaningful difference was observed between the mean of erectile dysfunction score and economic condition and smoking ( $p>0.05$ ).

The results of Pearson coefficient correlation test showed a significant relationship between erectile dysfunction in men with heart failure and age ( $p<0.001$ ), hemoglobin level ( $p=0.003$ ), left ventricular ejection fraction ( $p=0.030$ ), cholesterol ( $p=0.001$ ), depression ( $p<0.001$ ), and quality of life ( $p<0.001$ ) (Table. 1).

According to Table 2, t-test showed a significant difference between the mean of erectile dysfunction and renal disease ( $p=0.009$ ).

The following results were obtained: the mean of erectile dysfunction score based on the use of digoxin ( $p=0.014$ ), angiotensin receptor blockers ( $p<0.001$ ), beta blockers ( $p=0.001$ ), and diuretics ( $p=0.035$ ). However, there was no statistically significant difference based on the use of angiotensin converting enzyme inhibitors, nitrates, calcium blocker, and aldactone ( $p>0.05$ ).

## Discussion

In the present study, 80% of men with systolic heart failure suffered from erectile dysfunction and 36% of them had severe erectile dysfunction. This finding correlated with findings of other researchers. In a survey conducted by Medina, 74% [9], Schwartz, 84% [10] and Rastogi, 75% of patients with heart failure had erectile dysfunction [11]. The present findings support the claim that sexual aspect changes in patients with heart failure. Hence it is recommended that

Table 2: Comparison of the mean of erectile dysfunction score based on having morbidity in men with systolic heart failure

Having other morbidity	Erectile dysfunction
	Mean $\pm$ SD
Diabetes	+ 13.57 $\pm$ 6.05
	- 14.23 $\pm$ 6.39
Statistical test	T=-0.496 p=0.621
Hypertension	+ 15.45 $\pm$ 6.31
	- 13.31 $\pm$ 6.16
Statistical test	T=1.621 p=0.108
Renal disease	+ 8.14 $\pm$ 2.91
	- 14.46 $\pm$ 6.23
Statistical test	T=-2.652 p=0.009
Chronic obstructive pulmonary disease	+ 14.65 $\pm$ 6.89
	- 13.83 $\pm$ 6.09
Statistical test	T=0.550 p=.584
Ischemic heart disease	+ 14.66 $\pm$ 6.16
	- 13.81 $\pm$ 6.32
Statistical test	T=0.578 p=0.564
Asthma	+ 9.60 $\pm$ 4.77
	- 14.25 $\pm$ 6.26
Statistical test	T=-1.632 p=0.106

effective ways are considered to improve sexual function in these patients for promoting their quality of life.

The present study indicated a statistically significant relationship between age and intensity of erectile dysfunction; in a way that with an increase in age, intensity of erectile dysfunction was increased too. This finding is similar to the results of the studies done by Apostolo [16] and Stink [15]. They showed that there was a relationship between erectile dysfunction and age in men with heart failure. Using research evidence, many researchers proved the effect of age variable in erectile dysfunction [12, 13]. The results of Tels's study suggested that the incidence of sexual dysfunction in men aged from 40 to 49 years old was 29%, 50 to 59 years old was 50%, and 60 to 69 years old was 74%. In addition, the incidence of erectile dysfunction in men aged from 40 to 49 years old had increased by 1%, 50 to 59 years old by 2%, and 60 to 69 years by 10% [12]. Lindav's study suggested that an increase in age led to a decrease in sexual activity; in a way that 73% of men aged from 57 to 64 years old, 53% of men from 65 to 74 years old, and 26% of men from 75 to 85 years old were sexually active. In this study, 37% of the participants had erectile dysfunction and 4% of them used medicines to achieve an erection [13]. In Gerson's study, the age from 65 to 75 was one of the most important risk factors for erectile dysfunction; in a way that 62.3% of men between 65 and 75 and 21.6% of men between 45 and 55 suffered from erectile dysfunction [26]. Changes in penile vessels and blood circulation, and reduction of androgen, smooth muscles cell, and nitric oxide formation probably increase erectile dysfunction in patients with heart failure as they grow older.

The results of the present study showed that with reduction in left ventricular ejection fraction, intensity of erectile dysfunction in men with systolic heart failure would increase. This correlates with the findings of Mondress and Jaarsma's study in 1996. In their study, they indicated that reduction in left ventricular ejection fraction had a negative impact on sexual interest, sexual function and marital relationships [17, 27]. Reduction of left ventricular ejection fraction

probably limits penile vessels congestion possibility to achieve enough erection and as a result helps erectile dysfunction.

The present study suggested that a decrease in hemoglobin level is associated with an increase in the intensity of erectile dysfunction. This finding is similar to the results of Apostolo's study that showed a relationship between blood hemoglobin level and erectile dysfunction in men with heart failure [16]. Anemia may intensify sexual dysfunction by reducing tolerance level of activity and causing fatigue.

In the present study, there was a relationship between blood cholesterol level and erectile dysfunction. This finding correlates with the results of Vorntzez and Smith's study which showed a relationship between an increase in blood cholesterol level and erectile dysfunction [28, 29].

A systematic study done by Meiner indicated that dyslipidemia and hyperlipidemia increase the possibility of vascular disorders related to erectile dysfunction [30]. Therefore, perhaps lifestyle modification through nutrition, exercise and appropriate medication to control and decrease blood cholesterol can reduce the intensity of erectile dysfunction in men with heart failure.

There was a significant relationship between erectile dysfunction and education and occupation in the present study; in a way that the highest rate of erectile dysfunction was related to the uneducated and retired people. This finding is consistent with the results of Hameldon's study. He believes that demographic and social variables such as education and occupation are associated with sexual dysfunction [31].

In the present study, men with heart failure who also suffered from renal disease had more severe erectile dysfunction. This finding is consistent with the results of Jaarsma's study that showed chronic diseases together with heart failure is associated with an increase in the intensity of erectile dysfunction [6]. In Palmer's study, the incidence of sexual dysfunction in men with chronic renal failure undergoing hemodialysis was reported to be 86.6% [32]. Similarly, in

Nasir's study, 71% of men undergoing dialysis had erectile dysfunction [33]. It seems that mental and psychological factors, atherosclerosis process intensification in renal diseases, hormonal changes in the form of hyperprolactinemia and reduction in serum testosterone level, and side effects of controlling blood pressure medications increase erectile dysfunction in men with heart failure and renal function disorder.

In the present study, men with systolic heart failure who were treated with the drug digoxin had more erectile dysfunction. This is consistent with the findings of Rastogi's study which showed that the side effects of some drugs such as digoxin are related to erectile dysfunction in men with heart failure [11]. Schwartz claimed that the use of digoxin was associated with an increase in the incidence of erectile dysfunction; in a way that there was a relationship between an increase in estrogen level and a decrease in lutein hormones and testosterone and the use of digoxin. Digoxin also causes dysfunction in nitric oxide production and disorder in vascular smooth muscles [34].

In the present study, men with heart failure treated with angiotensin receptor blockers had more erectile dysfunction. This correlates with the findings of Rastogi's study [11]. Nevertheless, some studies have revealed that angiotensin receptor blockers have helped sexual function [34, 36].

In the present study, men with systolic heart failure who were treated with beta blockers had more erectile dysfunction. These findings are similar to the results of Rastogi's study [11]. Sylvestrey's study suggested that 31% of patients with cardiovascular problems had faced sexual disorders after starting treatment with atenolol due to its side effects [37]; however, Coroic Shank's study indicated that metoprolol and atenolol had no effect on sexual function [38].

In the present study, men with heart failure who were treated with diuretics had more erectile dysfunction. This finding correlates with the results of Apostolo's study which showed there was a relationship between the use of diuretics and erectile dysfunction [16].

Buranakit Jaron also found out that treatment with diuretic in men caused sexual dysfunction (CI=95% 1.3-2.8, OR=1.9). In general, side effects of medications can affect patients in following medicinal instructions. Since some medicines cause or intensify erectile dysfunction in men with heart failure, their negative effects on different aspects of patients' life, including sexual aspect, can influence on following medicinal instructions. The present study showed that men with systolic heart failure who had more severe erectile dysfunction suffered from a deeper depression. This is consistent with the results of other researchers who revealed that mood disorders and depression are associated with erectile dysfunction [34, 40, 41]. Like depression which is followed by a decrease in sexual preferences, mental and psychological factors also bring about sexual disorders in men with systolic heart failure.

The present study revealed that men with higher intensity of erectile dysfunction experience lower quality of life. This correlates with the results of the studies carried out by Schwartz [10] and Fritaz [42]. It seems that sexual function is an important aspect in the quality of life of patients with heart failure; in a way that its dysfunction reduces the quality of life and negatively affects interpersonal relationship, and mental and psychological aspects.

## Conclusion

In the present study, erectile dysfunction was common among patients with systolic heart failure and was related to factors such as age, occupation and education, left ventricular ejection fraction, hemoglobin and cholesterol level, renal disease, type of medications, depression and quality of life. The main limitation of this study was ignoring the control group. It is suggested that further studies are done to determine the frequency of erectile dysfunction in men with diastolic failure.

## Acknowledgments

This study is part of a research project with the code 1.11972. Hereby, we thank the

Deputy of Research in Islamic Azad University, Karaj branch, for approving the research project and his financial support.

## References

1. Zeighami Mohammadi Sh, Asgharzadeh Haghighi S, Nikbakht Nasrabadi A. The Study of Electrolyte Abnormalities and Renal Dysfunction on Elderly Patients with Systolic Heart Failure. *Iranian J Crit Care Nurs*. 2010;3(4):153-64.
2. Davis R, Hobbs F, Lip G. ABC of heart failure. *BMJ*. 2000;320:39-42.
3. Shojaei F. Quality of Life in Patients with Heart Failure. *Hayat*. 2009;14(2):5-13. [Persian]
4. Abbasi A, Asayesh H, Hosseini SA, Ghorbani M, Abdollah AA, Rouhi GH, et al. The relationship between functional performance in patients with heart failure and quality of life. *Iranian South Med J*. 2010;13(1):31-40. [Persian]
5. Rezaaei Looye H, Dalvandi A, Hosseini M, Rahgozar M. The Effect of Self Care Education on Quality of Life among Patients with Heart Failure. *J Rehabilitation*. 2009;10(2):21-6. [Persian]
6. Jaarsma T. Sexual problems in heart failure patients. *Eur J Cardiovasc Nurs*. 2002 Feb;1(1):61-7.
7. Schwarz ER, Rastogi S, Kapur V, Sulemanjee N, Rodriguez JJ. Erectile Dysfunction in Heart Failure Patients. *J Am Coll Cardiol*. 2006;48(6):1111-9.
8. Moeen MR. Male erectile dysfunction . Problems a new strategies management. *The Iranian Journal of Shahid Sadoughi Yazd University of Medical Sciences and Health Services*. 2002;10(2):95-105. [Persian]
9. Medina M, Walker C, Steinke EE, Wright DW, Mosack V, Farhoud MH. Sexual concerns and sexual counseling in heart failure. *Prog Cardiovasc Nurs*. 2009;24(4):141-8.
10. Schwarz ER, Kapur V, Bionat S, Rastogi S, Gupta R, Rosanio S. The prevalence and clinical relevance of sexual dysfunction in women and men with chronic heart failure. *Int J Impot Res*. 2008;20(1):85-91.
11. Rastogi S, Rodriguez JJ, Kapur V, Schwarz ER. Why do patients with heart failure suffer from erectile dysfunction? A critical review and suggestions on how to approach this problem. *Int J Impot Res*. 2005;17(1):25-36.
12. Teles AG, Carreira M, Alarco V, Sociol D, Arag JM, Lopes L, et al. Prevalence, severity, and risk factors for erectile dysfunction in a representative sample of 3/548 portuguese men aged 40 to 69 years attending primary healthcare centers: results of the portuguese erectile dysfunction study. *J Sex Med*. 2008;5(6):1317-24.
13. Lindau ST, Schumm LP, Laumann EO, Levinson W, O'Muircheartaigh CA, Waite LJ. A study of sexuality and health among older adults in the United States. *N Engl J Med*. 2007;357(8):762-74.
14. Rahman MM, Laher I. Structural and functional alteration of blood vessels caused by cigarette smoking: an overview of molecular mechanisms. *Curr Vasc Pharmacol*. 2007;5:276-92.
15. Steinke EE, Mosack V, Wright DW, Chung ML, Moser DK. Risk factors as predictors of sexual activity in heart failure. *Dimens Crit Care Nurs*. 2009;28(3):123-9.
16. Apostolo A, Vignati C, Brusoni D, Cattadori G, Contini M, Veglia F, et al. Erectile dysfunction in heart failure: correlation with severity, exercise performance, comorbidities, and heart failure treatment. *J Sex Med*. 2009;6(10):2795-805.
17. Mandras SA, Uber PA, Mehra MR. Sexual activity and chronic heart failure. *Mayo Clin Proc*. 2007;82:1203-10.
18. Baumhäkel M, Böhm M. Erectile dysfunction correlates with left ventricular function and precedes cardiovascular events in cardiovascular high-risk patients. *Int J Clin Pract*. 2007;61(3):361-6.
19. Mehrabi F, Ehssanmanesh M, Karimi keisimi E. Demographic characteristics of men with sexual dysfunction, *Iranian Journal of Psychiatry and Clinical Psycholog*. 2003;9:14-22. [Persian]
20. Asadi Far F. The Attitudes to sexual dysfunction. *Tehran: Bersat*; 2001. [Persian]
21. Bener A, Al-Ansari A, Al-Hamaq AO, Elbagi IE, Afifi M. Prevalence of erectile dysfunction among hypertensive and nonhypertensive Qatari men. *Medicina (Kaunas)*. 2007;43(11):870-8.
22. Abbasi A, Nasiri H. Specific instrument for assessment quality of life of chronic heart failure patients. *Journal of Gorgan Bouyeh Faculty Nursing and Midwifery*. 2006;3(2):41-5. [Persian]
23. Kantor J, Bilker WB, Glasser DW, Margolis DJ. Prevalence of Erectile Dysfunction and Active Depression: An Analytic Cross-Sectional Study of General Medical Patients. *Am J Epidemiol*. 2002;156(11):1035-42.
24. Sajadi M, Akbari A, Kianmehr M, Ataroudi A. The relationship between self care and depression in patients undergoing hemodialysis. *Ofoh-E-Danesh*. 2008;14(1):13-7. [Persian]
25. Clark CM, Mahoney JS, Clark DJ, Eriksen LR. Screening for depression in Hepatitis C population: the reliability and validity of the Center for Epidemiologic Studies Depression Scale (CES-D). *J Adv Nurs*. 2002;40(3):361-9.
26. Geirsson G, Thornorgeirsson G, Guethmundsson O, Einarsson G. Risk factors and prevalence of erectile dysfunction amongst Icelandic men aged 45-75. *Laeknabladid*. 2006;92(7-8):533-7.
27. Jaarsma T, Dracup K, Walden J, Stevenson LW. Sexual function in patients with advanced heart failure. *Heart Lung*. 1996;25(4):262-70.
28. Vrentzos GE, Paraskevas KI, Mikhailidis DP. Dyslipidemia as a risk factor for erectile

- dysfunction. *Curr Med Chem.* 2007;14(16):1765-70.
29. Smith NJ, Sak SC, Baldo O, Eardley I. The prevalence of newly diagnosed hyperlipidaemia in men with erectile dysfunction. *BJU Int.* 2007;100(2):357-61.
  30. Miner M, Billups KL. Erectile dysfunction and dyslipidemia: relevance and role of phosphodiesterase type-5 inhibitors and statins. *J Sex Med.* 2008;5(5):1066-78.
  31. Holden CA, McLachlan RI, Pitts M, Cumming R, Wittert G, Ehsani JP, et al. Determinants of male reproductive health disorders: the Men in Australia Telephone Survey (MATeS). *BMC Public Health.* 2010;10:96.
  32. Makarem AR, Karami MY, Zekavat OR. Erectile dysfunction among hemodialysis patients. *Int Urol Nephrol.* 2011;43(1):117-23.
  33. Nassir A. Erectile dysfunction risk factors for patients entering dialysis programme. *Andrologia.* 2010;42(1):41-7.
  34. Doumas M, Douma S. The effect of antihypertensive drugs on erectile function: a proposed management algorithm. *J Clin Hypertens (Greenwich).* 2006;8(5):359-64.
  35. Shindel AW, Kishore S, Lue TF. Drugs designed to improve endothelial function: effects on erectile dysfunction. *Curr Pharm.* 2008;14(35):3758-67.
  36. Franzen D, Metha A, Seifert N, Braun M, Hopp HW. Effects of beta-blockers on sexual performance in men with coronary heart disease: A prospective randomized and double blinded study. *Int J Impot Res.* 2001;13(6):348-51.
  37. Cruickshank JM. Beta-blockers and heart failure. *Indian Heart J.* 2010;62(2):101-10.
  38. Buranakitjaroen P. Prevalence of Erectile Dysfunction among Treated Hypertensive Males. *J Med Assoc Thai.* 2006;89 Suppl 5:S28-36.
  39. Makhoulouf A, Kparker A, Niederberger CS. Depression and erectile dysfunction. *Urol Clin North Am.* 2007;34(4):565-74, vii.
  40. Zemishlany Z, Weizman A. The impact of mental illness on sexual dysfunction. *Adv Psychosom Med.* 2008;29:89-106.
  41. Freitas D, Athanazio R, Almeida D, Dantas N, Reis F. Sildenafil improves quality of life in men with heart failure and erectile dysfunction. *Int J Impot Res.* 2006;18(2):210-2.