



Investigating the effects of Benson's relaxation technique on quality of life among patients receiving hemodialysis

Hossein Feyzi¹, Behnam Khaledi Paveh^{*1}, Fatemeh Hadadian¹, Mansoor Rezaie², Mojtaba Ahmadi³

^{*1}.Department of Nursing, School of Nursing and Midwifery, Kermanshah University of Medical Sciences, Kermanshah, Iran

². Department of Biostatistics and Epidemiology, Center for Social Development and Health Promotion, University of Medical Sciences, Kermanshah, Kermanshah, Iran

³. Department of Psychology, School of Medicine, Kermanshah University of Medical Sciences, Kermanshah, Iran

ARTICLE INFO

Article type:
Original article

Article history:
Received: 26 Apr 2014
Revised: 10 Feb 2015
Accepted: 12 Feb 2015

Keywords:
Hemodialysis
Quality of life,
Benson's relaxation technique

ABSTRACT

Aims: End-stage renal disease is a major health challenge worldwide. Patients with this disease need to receive frequent hemodialysis. The disease and hemodialysis both negatively affect patients' quality of life. The aim of this study was to investigate the effects of Benson's relaxation technique on quality of life among patients receiving hemodialysis.

Methods: A convenience sample of 60 patients receiving hemodialysis in Imam Reza Hospital, Kermanshah, Iran, was recruited to this clinical trial. Initially, patients' quality of life was evaluated by a disease-specific questionnaire. Then, patients were randomly allocated to either the control or the treatment groups. Patients in the treatment group were trained to perform Benson's relaxation technique at their home on a daily basis for 60 consecutive days. Patients in the control group received no relaxation intervention. Immediately after the study, the same questionnaire was used for reevaluating patients' quality of life. The independent-samples t-test was used for data analysis.

Results: After the study, the mean scores of general and total quality of life were significantly higher than the corresponding pretest readings ($p = 0.001$ and 0.02 , respectively). However, the mean score of specific quality of life did not change significantly after the study ($p=0.2$).

Conclusions: Benson's relaxation technique has positive effects on quality of life among patients receiving hemodialysis and therefore, it can be used for enhancing patients' quality of life in hemodialysis units.

Please cite this paper as:

Feyzi H, Khaledi Paveh B, Hadadian F, Rezaie M, Ahmadi M . Investigating the effects of Benson's relaxation technique on quality of life among patients receiving hemodialysis. Iran J Crit Care Nurs. 2015;8(1):13-20.

1. Introduction

Currently, one of the major healthcare problems worldwide is the increasing

prevalence of chronic diseases [1]. A chronic disease is a medical condition or a health problem which is associated with different symptoms and disabilities and needs long-term management [2]. One of the commonest chronic diseases is end-stage renal disease

*Correspondence Author: Behnam Khaledi Paveh
Department of Psychiatry, School of Medicine, Kermanshah
University of Medical Sciences, Kermanshah, Iran. Tel:+98-
9181731736
Email: bkhaledi@yahoo.com

(ESRD). ESRD is a progressive and irreversible kidney disorder in which the body fails to maintain fluid and electrolytes balance and the afflicted patient may finally develop uremia or azotemia [3]. Hence, it is a major health problem throughout the world [4].

Statistics show that the number of patients with ESRD worldwide is more than two million from which, 1.38 million patients receive hemodialysis and 170000 ones undergo peritoneal dialysis [5]. Currently, there are 450000 patients with ESRD in the United States [6]. This number will increase to 2.24 million patients by the year 2030 [7]. In our country, Iran, the incidence of ESRD is around 22.6%. Consequently, 4000 new cases of ESRD are diagnosed each year. Currently, the number of patients with ESRD in Iran is greater than 33000 [8].

ESRD and its complications significantly affect afflicted patients' lifestyle, health and well-being, social and familial roles, and quality of life (QOL). One of the reasons for initiating hemodialysis for patients with ESRD is its positive effects on patients' sense of well-being [9]. However, although dialysis significantly improves survival, QOL of patients who receive dialysis is lower than healthy people [10]. Valderrabano et al. (2001) noted that dialysis and other treatments alleviate the symptoms of ESRD and promotes patients' lifestyle; nonetheless, patients' QOL is compromised due the complications of ESRD and most of these patients experience different degrees of disability [11]. Feroze et al. (2011) also reported that QOL among patients receiving hemodialysis is poor [12].

Poor QOL affects all aspects of patients' lives. According to Rambod et al. (2008), hemodialysis negatively affects patients' functional status and impairs their ability to perform activities of daily living [13]. Moreover, as these patients need to receive frequent dialysis treatments, they may be either unable to find a regular and continued employment or lose their employment and therefore, experience financial problems [14].

In addition, ESRD- and dialysis-related fatigue, anxiety, and neurohormonal disorders negatively affect psychological, familial, and social aspects of patients' lives [15] and hence, compromise the social aspect of QOL [16]. Mehraban et al. (2011) found that the QOL of patients receiving kidney transplantation and peritoneal hemodialysis is similar to that of healthy people and significantly higher than patients undergoing hemodialysis [17]. However, Smaeli et al. (2005) reported a good QOL among Iranian patients receiving hemodialysis [18].

Most Americans use complementary and alternative modalities for promoting their health and recovering from diseases. Studies showed that about 50% of Americans use these modalities as part of their treatment plan. The use of complementary and alternative modalities is increasing progressively. Complementary and alternative modalities include a wide range of modalities such as Chinese medicine, yoga, acupuncture, homeopathy, naturopathy, music therapy, energy healing approaches, chiropractic, body mind interaction, and relaxation [19].

One of the complementary and alternative modalities is Benson's relaxation technique. Herbert Benson based his technique on the concepts of transcendental meditation. In this technique, a person adopts a passive mode and allows relaxation to arise and progress [20]. The progressive muscle relaxation technique (PMR) was developed for alleviating tension and anxiety. The aim of this cost-effective, non-invasive, self-administered technique is to enhance people's awareness of muscular tension and relaxation and also to help them relax their muscles [21]. Shariati et al. (1982) found that PMR was effective in relieving anxiety among patients receiving hemodialysis [22]. Yildirim and Fadiloglu (2006) also found that PMR significantly alleviated state anxiety and enhanced QOL among patients undergoing dialysis [23]. Studies conducted in Iran also have shown the positive effects of PMR on QOL among patients with breast cancer [24]

and rheumatoid arthritis [25] as well as patients undergoing coronary artery bypass graft [26].

To the best of our knowledge, no studies in Iran have yet investigated the effects of muscle relaxation on QOL among patients undergoing hemodialysis. Accordingly, this study was conducted to provide evidence in this area. The aim of the study was to investigate the effects of Benson's relaxation technique on QOL among patients receiving hemodialysis.

2. Methods

This was a quasi-experimental clinical trial. The population of the study consisted of all patients receiving hemodialysis in Kermanshah, Iran. A convenience sample of 60 patients receiving hemodialysis in Imam Reza Hospital, Kermanshah, Iran, was recruited. Patients were randomly assigned to either the control or the treatment groups. Patients were included in the study if they were desired for participation, had no debilitating underlying condition which interfered with implementing relaxation, were able to establish face-to-face communication, and neither performed relaxation or stretching exercise at the time of the study nor used narcotics. The exclusion criteria were patients' withdrawing from the study, using tranquilizers, anxiolytic agents, or antidepressants, requiring hospitalization, failing to perform three consecutive or five intermittent relaxation sessions, and using treatments other than hemodialysis (such as peritoneal dialysis or kidney transplantation).

A demographic questionnaire and the Kidney Disease Quality of Life Short Form (KDQOL-SF) were used for data collection. The KDQOL-SF is a disease-specific questionnaire for assessing QOL among patients undergoing hemodialysis and consists of two main domains including general QOL and specific QOL. The general domain includes two physical and emotional sub-domains. The physical sub-domain, in turn, consists of four sub-scales including general health (six items), physical functioning (ten items), physical role

performance (four items), and physical pain (three items) while the emotional sub-domain includes three sub-scales of emotional role performance (three items), social functioning (two items), and mental health (eight items). On the other hand, the specific domain of the KDQOL-SF consists of nine sub-scales including limitations related to kidney disease (eleven items), health-related mental problems (six items), health-related physical performance (twelve items), general health status (three items), health-related family satisfaction (four items), sleep status (on a 0–100 rating scale), health-related employment status (three items), sexual problems (two items), and satisfaction with both care and healthcare professionals (three items). The total score of each sub-scale ranges from 0 to 100. Moreover, the total scores of the two domains of QOL were classified as poor (0–33), moderate (34–66), and good (> 66). The reliability and the content validity of the Persian version of KDQOL-SF had been evaluated and confirmed by Rahimi et al (2006) [27]. They used the split-half technique and reported a correlation coefficient of 0.9 for the questionnaire, confirming its reliability.

Initially, eligible patients were informed about the aim and the methods of the study and were asked to sign the study informed consent form. Then, after measuring their QOL by using the KDQOL-SF, patients were allocated to either the control or the treatment groups. Patients in the treatment group were trained to implement Benson's relaxation technique. Trainings were continued until patients were able to independently perform relaxation. All trainings were provided in one session. In the next day following trainings, patients were provided with an audio file and a written booklet containing trainings about Benson's relaxation technique. Patients were asked to perform the relaxation exercises at home on a daily basis for two months—60 sessions in total. The six steps of Benson's relaxation technique were as follows:

1. Assume a comfortable position;

Table 1: The mean scores of the general QOL sub-scales before and after the study

The sub-scales of the general QOL domain		Mean	Standard deviation	Z	P
General health	Before	56.57	17.61	- 2.02	0.04
	After	63.60	18.90		
Physical functioning	Before	56.96	23	- 2.36	0.01
	After	70.03	23.61		
Physical role performance	Before	39.86	24.55	- 1.80	0.07
	After	49.82	25.82		
Emotional role performance	Before	61.48	23.29	- 2.20	0.04
	After	73.74	19		
Social functioning	Before	44.06	24.03	- 4.12	< 0.001
	After	69.16	33.45		
Physical pain	Before	52.43	31.47	- 1.89	0.05
	After	61.96	29.52		
Mental health	Before	59.65	24.61	- 1.72	0.08
	After	66.10	21.68		

2. Close the eyes gently;
3. Deeply relax all the muscles of your body. Start from the sole and gradually move towards face. Keep muscles relaxed.
4. Breathe normally through your nose. Try to become aware of your breathing and say a single word softly at each exhalation.
5. Continue this exercise for 15–20 minutes. Try to keep all your muscles relaxed. After 15–20 minutes, open your eyes gently and remain in place for a few minutes.
6. Avoid being worried about whether having relaxation or not. Instead, let relaxation happens gradually. Push away disturbing thoughts.

The SPSS16 was used for conducting the Mann-Whitney as well as the paired- and the independent-samples t tests.

Official permissions for conducting the study were obtained from Kermanshah Faculty of Nursing and Midwifery, the Research council of Kermanshah University of Medical Sciences, and the administrators of Imam Reza Hospital, Kermanshah, Iran. Then, the researchers and the research project were introduced to the participants and a written consent was obtained from them. Research

participants were ensured that their information will be managed and reported confidentially. They were completely free to either participate or reject participation in the study.

3. Results

In total, 60 patients participated in the study from which, 57% were female. The mean, the minimum, and the maximum of participants' ages were 45.31 ± 14.70 , seventeen, and 75 years, respectively. Most participants were single (58%) and housewives (50%), held secondary diploma (33%), and had sufficient income (51%). The means of the length of having kidney disease and receiving hemodialysis were 5.48 ± 6.10 and 3.19 ± 2.96 years, respectively.

About 93% of the participants knew about their kidney disease and strategies for its management while 88% of them reported that they required additional information about their disease.

Disease-related limitations were tolerable, almost tolerable, and intolerable for respectively 33%, 54%, and 13% of the participants.

Table 2: The mean scores of the specific QOL sub-scales before and after the study

The sub-scales of the specific QOL domain		Mean	Standard deviation	Z	P
Disease-related limitations	Before	45.17	13.79	- 1.16	0.24
	After	50.47	13.16		
Mental health	Before	5.28	10.09	- 0.73	0.46
	After	56.71	8.91		
Physical functioning	Before	66.82	17.52	- 2.69	0.007
	After	76.10	17.68		
General health status	Before	53.60	28.39	- 0.58	0.55
	After	55.82	23.99		
family satisfaction	Before	63.23	25.82	- 2.08	0.03
	After	73.70	27.27		
Sleep status	Before	9.89	17.47	- 0.98	0.32
	After	7.44	1.99		
Employment status	Before	8.64	1.90	- 0.65	0.51
	After	8.39	1.68		
Sexual problems	Before	50	43.73	- 0.05	0.95
	After	51.17	36.03		
Satisfaction with healthcare professionals	Before	8.23	1.43	- 0.02	0.98
	After	8.26	1.61		

Study findings revealed that in the treatment group, except for the physical role performance ($p=0.07$) and the mental health ($p=0.08$) subscales, the posttest values of the other five sub-scales of general QOL (i.e. general health, physical functioning, physical pain, emotional role performance, and social functioning) were significantly higher than the corresponding pretest values ($p<0.05$; Table 1). On the other hand, only the posttest readings of the health-related physical functioning ($p=0.007$) and the health-related family satisfaction ($p=0.03$) sub-scales of the specific domain of QOL were significantly higher than their pretest readings ($p<0.05$).

Moreover, in the treatment group, the mean

scores of the general domain of QOL ($p=0.001$) and total QOL ($p=0.01$) significantly increased after the study while the changes in the mean score of the specific domain of QOL were statistically insignificant ($p=0.2$; Table 3).

Before the intervention, the general QOL of only 12.3% of participants was good.

This value increased to 30.2% after the study. Moreover, before the intervention, 38.7% and 61.3% of participants had respectively poor and moderate specific QOL while after the intervention, these values changed to 28.1% and 71.9%, respectively.

Finally, before the study, the total QOL of 20% and 80% of the participants was respectively poor and moderate; meanwhile,

Table 3: The mean scores of specific, general, and total QOL in both groups before and after the study

The domains of QOL		Mean and standard deviation	T	P
General QOL	Before	- 2.53±14.89	- 3.64	0.001
	After	13.72±14.68		
Specific QOL	Before	4.03±9.08	- 1.21	0.2
	After	0.06±6.47		
Total QOL	Before	0.66±10.44	- 2.35	0.01
	After	10.48±11.15		

Table 4: The mean scores of different levels of specific, general, and total QOL in both groups before and after the study

Total QOL and its domains	Levels	Before	After
The general QOL domain	Poor (0–33)	20.4%	15.1%
	Moderate (34–66)	67.3%	54.7%
	Good (> 66)	12.3%	30.2%
The specific QOL domain	Poor (0–33)	38.7%	28.1%
	Moderate (34–66)	61.3	71.9%
Total QOL	Poor (0–33)	20%	12.5%
	Moderate (34–66)	80%	84.4%
	Good (> 66)	---	3.1%

these values changed to 12.5% and 84.4%, respectively after the study and the total QOL of 3.1% participants turned good (Table 4).

4. Discussion

This study was conducted to investigate the effects of Benson's relaxation technique on QOL among patients receiving hemodialysis. The findings of the study indicated the effectiveness of Benson's relaxation technique in enhancing patients' QOL.

After the study, the mean scores of different sub-scales of the general QOL were significantly higher than the pretest readings, confirming the positive effects of Benson's relaxation technique on QOL. These findings are in line with the findings reported by Shariati et al. (2011) [22].

They found that Benson's relaxation technique significantly improved all aspects of QOL, except for the sexual activity sub-scale, among patients with breast cancer [24]. Our findings revealed that Benson's relaxation technique significantly alleviated patients' physical pain. Fayazi et al. (2010) also found that this relaxation technique was effective in relieving postoperative pain among patients undergoing coronary artery bypass graft [29].

Moreover, Shaban et al. (2006) found that both progressive muscle relaxation and music therapy significantly reduced pain among patients with cancer [30].

Dehdari et al. (2009) also reported that progressive muscle relaxation was effective in

improving all aspects of QOL and alleviating state and trait anxiety after coronary artery bypass graft surgery [24]. However, our intervention had no significant effect on the mental health sub-scale of QOL. This conflicting finding can be attributed to the difference in the samples of the two studies. ESRD is a chronic condition which affects QOL.

Moreover, hemodialysis-related complications also reduce the QOL of patients with ESRD. Accordingly, the combined negative effects of ESRD and hemodialysis on QOL may be much greater than the effects of a short-term coronary artery bypass graft.

We also found that apart from the physical functioning and the health-related family satisfaction sub-scales, the Benson's relaxation technique had no significant effect on the other sub-scales of the specific domain of QOL. However, Ashktorab et al. (2012) reported that progressive muscle relaxation significantly affected sleep quality among patients undergoing hemodialysis.

This contradiction between the findings of the two studies is probably due to the differences in the instruments and the interventions of the studies.

Ashktorab et al. (2012) assessed only patients' sleep quality by using the Pittsburgh Sleep Quality Index while in our study, sleep quality was evaluated as a sub-scale of QOL. Moreover, we used Benson's relaxation technique while the intervention implemented

by Ashktorab et al. (2012) was Jacobson's relaxation technique.

Study findings also revealed that Benson's relaxation technique significantly improved patients' general and total QOL while having no significant effect on their specific QOL. Masoudi et al. (2011) also reported that progressive muscle relaxation was effective in enhancing QOL among caregivers of patients with multiple sclerosis [31].

Moreover, Yildirim and Fadiloglu (2006) found that progressive muscle relaxation significantly alleviated state and trait anxiety and enhanced QOL in patients receiving dialysis [23]. Zamanzadeh (2008) also reported the same findings [32].

5. Conclusions

The Benson's relaxation technique has positive effects on QOL among patients receiving hemodialysis. Accordingly, this technique can be used for enhancing patients' QOL in hemodialysis units.

6. Acknowledgments

We would like to sincerely thank all patients who participated in this study. In addition, we are heartily grateful to the staffs and the head-nurse of the dialysis unit of Imam Reza Hospital as well as the Research Councils of Kermanshah Faculty of nursing and Midwifery and Kermanshah University of Medical Sciences, Kermanshah, Iran.

References

1. Kasper D. Principles of internal medicine. 16thed, Mcm raw Hill Medical. Newyork. 2005.
2. Nobahar M, Baba Mohamadi H, Soleymani M, Asgari MR, Vafiie A. Comprehensive text book of medical- surgical nursing. 2st ed. Tehran; Boshra 2008. [Persian]
3. Tayebi A, Raisifar A, Ebadi A, Eynolahi A, Eynolahi B, Rafeian Z, Keyvanlo F. Review the renal transplantation patients' quality of lifeby using kidney transplantation questionnaire. Iranian Journal of Critical Care Nursing. 2010;3(3):125-28. [Persian]
4. Rahimi A, Ahmadi F, GHaliaf M. Effect of continuous care model on stress, anxiety and depression in hemodialysis patients. Journal of Medicine. 2006;30(4):353-95. [Persian]
5. Hojat M. Efficiency hemodialysis in the chronic kidney disease. Iranian Journal of Critical Care Nursing. 2009;2(2):61-6. [Persian]
6. Harrison's Internal Medicine. Hasan Mirzaie B, Ebrahimipour H. 2st ed. Tehran; Samat- Andishe Rafie. 2010;133-38. [Persian translator]
7. Lo DS, Rabbat CG, Clase CM. Thromboembolism and anticoagulant management in hemodialysis patients: A practical guide to clinical management. Thrombosis Research, Elsevier Ltd. 2006;118(3):385-95.
8. Rahimzadeh A. Anticoagulation. In: Mohsenfar. Hemodialysis Efficiency. Center Manegment Transplantation and Spesific Disease/ Nurse and Hemodialysis. Lahzeh. 2006;91-101,123-24.
9. Kutner NG. Quality of life and daily hemodialysis. Seminars in dialysis. 2004;17(2):92-8.
10. Fukuhara S, Lopes AA, Bragg-Gresham JL, Kurokawa K, Mapes DL, Akizawa T L, et al. Heath related quality of life amongdialysis patient on three continents, the dialysis outcomes and practice patterns study. Kidney int. 2003;64:1903-10.
11. Valderrabano F, Jofre R, Lopez-Gomez JM. Quality of life in end-stage renal disease patients. Am J Kidney Dis. 2001;38(3):443-64.
12. Feroze U, Noori N, Kovesdy CP, Molnar MZ, Martin DJ, Reina-Patton A, et al. Quality of Life and Mortality in Hemodialysis Patients: Roles of Race and Nutritional Status. Clin J Am Soc Nephrol. 2011;6(5):1100-11.
13. Rambood M, Rafie F, Hosseyni F. Quality of life in patients with chronic renal failure. Journal of Nursing and Midwifery, Tehran University of Medical Sciences. 2008;14(2):51-61. [Persian]
14. Thomas CJ. The impact of religiosity, social support and health locus of control on the health related quality of life of African-American Hemodialysis patients. Ph. Dissertation, Norfolk Virginia, Norfolk State University. 2003.
15. Kimmel PL. Psychosocial factors in adult end-stage renal disease patients treated with hemodialysis: correlates and outcomes. Am J Kidney Dis. 2000;35(4):132-40.
16. Al-Arabi S. Quality of life: subjective descriptions of challenges to patients with end stage renal disease. Nephrology Nursing Journal. 2006;33(3):285-98.
17. Mehraban D, Salehi M, Naderi GH, Najafi I, Monirolsadat H. Compare the quality of life in patients with renal replacement therapy population. Journal of Health Management. 2005;8(20):68-76.
18. Smaeli M, Alikhani M, Kholam Eraghi M. The quality of life and self-efficacy of the patients

- under hemodialysis. *Iranian Journal of Nursing*. 2005; 18(41,42): 76-84.
19. Taylor C. Fundamentals of nursing. Azizi D, Taghafi A. Khadem Hosseini H. 1st ed. Tehran; Hayyan- Abasaleh Publication. 2008:718-30. [Persian]
 20. Renato D. Alarcon, Charlotte S. Jenkins, Diane E. Heestand, Lowell K. Scott, Larry Cantor. The effectiveness of progressive relaxation in chronic hemodialysis patients. *Journal of Chronic Diseases*. 1982;35(10):797-802.
 21. Yildirim YK, Fadiloglu C. The effect of progressive muscle relaxation training on anxiety levels and quality of life in dialysis patients. *Edtna Erca J*. 2006;32(2):86-8.
 22. Shariati AA, Salehi M, Ansari M, Latifi SM. Survey the Effect of Benson Relaxation Intervention on Quality of Life (QOL) in Breast Cancer Patients Undergoing Chemotherapy. *Sci Med J*. 2011;9(6):625-32. [Persian]
 23. Bagheri M, Mohseni MA, Shayesteh M.. The effect of Benson Relaxation Technique on rheumatoid arthritis patients: Extended report. *International Journal of Nursing Practice*. 2006;12:214-9. [Persian]
 24. Dehdari T, Heidarnia A, ramezankhani A, Sadeghian S, Ghofranipour F. Effects of progressive muscular relaxation training on quality of life in anxious patients after coronary artery bypass graft surgery. *Indian J Med Res*. 2009;129:603-8. [Persian]
 25. Kaplan S. Synopsis of psychiatry. Behavioral sciences/ Clinical psychiatry. Rafiei H. (Persian translator). 5th ed. Vol 2. Tehran; Arjmand publication. 2008.
 26. Ashktorab T, Saeedi M, Saatchi K, Zayeri F, Amir Ali Akbari S. The Effect of Progressive Muscle Relaxation on Sleep Quality of Patients Undergoing Hemodialysis. *Iranian Journal of Critical Care Nursing*. 2012;5(1):23-8. [Persian]
 27. Rahimi A, Ahmadi F, GHaliyaf M. Effect of continuous care model on quality of life in hemodialysis patients. *Iranian Journal of Medical Sciences*. 2006;13(52):354-59. [Persian]
 28. Ghafari S, Ahmadi F, Nabavi S, Memarian R. Effects of progressive muscle relaxation technique on depression, anxiety and stress in patients with multiple sclerosis. *J Med*. 2008;32(1):45-53. [Persian]
 29. Fayazi S, Shariati A, Momeni M, Latifi M. The efficacy of Benson's relaxation technique on postoperative pain in coronary artery bypass graft. *Sci Med J*. 2010;8(4):479-89. [Persian]
 30. Shaban M, Rasoolzadeh N, Mehran A, Moradalizadeh F. Effect of non-pharmacological techniques (progressive muscle relaxation and music therapy) on pain in cancer patients. *Journal of Nursing & Midwifery, Tehran University of Medical Sciences (Hayat)*. 2006;12(3):63-72. [Persian]
 31. Masoudi R, Soleimani MA, Moghadasi J, Qorbani M, Mehralian H, Bahrami N. Effect of progressive muscle relaxation program on self-efficacy and quality of life in caregivers of patients with multiple sclerosis. *JQUMS*. 2011;15(2):41-7. [Persian]
 32. Rambod M, Sharifi F, Pourali Mohammadi N, Pasyar N, Raffi F. Evaluation of the effect of Benson's relaxation technique on pain and quality of life of haemodialysis patients: a randomized controlled trial. *Int J Nurs Stud*. 2014;51(7): 964-73.
 33. Zamanzadeh V. The effect of exercise on quality of life in hemodialysis patients. *Journal of Tabriz University of Medical Sciences*. 2008;1(3):51-5. [Persian]