



## The effects of the family-centered orientation program on satisfaction with healthcare services among patients with coronary artery disease

Vahideh Karimi<sup>1</sup>, Nasrin Hanifi<sup>2\*</sup>, Nasrin Bahraminejad<sup>1</sup>, Soghrat Faghihzadeh<sup>3</sup>

1. Medical-Surgical Department, Faculty of Nursing and Midwifery, Zanzan University of Medical Sciences (ZUMS), Zanzan, Iran

\*2. Department of Critical Care Nursing, Faculty of Nursing and Midwifery, Zanzan University of Medical Sciences (ZUMS), Zanzan, Iran

3. Biostatistics and Epidemiology Department, Zanzan University of Medical Sciences (ZUMS), Zanzan, Iran

### ARTICLE INFO

Article type:  
Original article

Article history:  
Received: 14 Dec 2014  
Revised: 18 Apr 2015  
Accepted: 26 Sep 2015

Keywords:  
Orientation program  
Family-centered education  
Patient satisfaction  
Coronary artery disease  
Coronary care unit

### ABSTRACT

**Aims:** Hospitalization in critical care units is associated with anxiety for patients and family members due to encountering unfamiliar and strange equipment and procedures. Such anxiety and unfamiliarity can bring patients dissatisfaction. This study was conducted "to examine the effects of the Family-Centered Orientation Program on satisfaction with healthcare services among patients with coronary artery disease".

**Methods:** A convenience sample of 80 patients with coronary artery disease was recruited to this clinical trial study from the coronary care unit of a teaching hospital affiliated to Zanzan University of Medical Sciences, Zanzan, Iran. Patients were randomly allocated to the control and the experimental groups. The Family-Centered Orientation Program was implemented for patients in the experimental group. Patient satisfaction with care services was evaluated at the time of hospital discharge by using the Patient Satisfaction Scale. Study data were analyzed by employing the SPSS<sub>17</sub> software as well as the Kolmogorov-Smirnov, the independent-samples t, and the Fisher's exact tests.

**Results:** After the study, the level of satisfaction with care services in the experimental group was significantly higher than the control group ( $p < 0.0001$ ).

**Conclusion:** Family-Centered Orientation Program is effective in enhancing patient satisfaction. Using this program for improving care quality and enhancing patient satisfaction is recommended.

**Conclusions:** Conclusion: Family-Centered Orientation Program is effective in enhancing patient satisfaction. Using this program for improving care quality and enhancing patient satisfaction is recommended.

Please cite this paper as:

Karimi V, Hanifi N, Bahraminejad N, Faghihzadeh S. The effects of the family-centered orientation program on satisfaction with healthcare services among patients with coronary artery disease. Iran J Crit Care Nurs. 2015;8(3):143-148.

### 1. Introduction

Patients who are hospitalized in critical care units experience high levels of anxiety [1]. Beside life-threatening conditions, other factors such as hospitalization, unfamiliar environment, inability to communicate, mobility restrictions,

financial problems, loss of control, separation from family members, noisy and strange equipment, frequent diagnostic and treatment procedures, and sudden changes in household and professional responsibilities are among the most important stressors in critical care units [1

\* Correspondence Author: Nasrin Hanifi, Department of Critical Care Nursing, Faculty of Nursing and Midwifery, Zanzan University of Medical Science(ZUMS), Zanzan, Iran  
Tel:+9824-33772514, Email:nasrinhanifi@zums.ac.ir

and 2]. These stressors affect both patients and their family members [2–4].

In order to cope with their anxiety, patients and their families need timely and precise information regarding their concerns particularly during the first day of hospitalization. Such information should be provided by critical care nurses [5]. Fulfillment of patients and families' educational needs by nurses can alleviate their stress and anxiety [6–8]. However, a great deal of nurses' time is spent on managing patients' physical problems and hence, they fail to assess and alleviate patients and families' concerns [3].

One of the strategies for providing information to patients is the Family-Centered Orientation Program (FCOP) [5–9]. The FCOP can alleviate families' psychological stress, improve their perception of patients' underlying conditions and prognoses, facilitate their coping with changes, enhance the quality of decision making, increase the effectiveness of treatments, and promote patients' well-being [3, 5–9]. It enables nurses to establish effective communication with patients and families, provide informational and social support to them, and enhance their satisfaction with care services [3, 10, and 11]. Greater satisfaction with services fosters positive attitudes toward the related service-providing institutes [11–13] and increases customers' interest in referring to those institutes [14]. On the other hand, inadequate information provision to patients and their families about the environment, procedures, and equipment of critical care units can negatively affect their satisfaction with care services [3].

Patient satisfaction is one the criteria for care quality assurance [10–15]. Consequently, nurses need to adopt strategies for alleviating patients and families' post-hospitalization anxiety and thereby enhancing their satisfaction with care services.

Chan et al. (2012) found that orientation programs can significantly alleviate patients' stress and enhance their knowledge and satisfaction [16]. Mohammadi et al. (2012) also

found that a motivational program developed based on the expectancy theory significantly enhanced patient satisfaction [15]. Moreover, Dehghannayyeri and Aghajani (2007) reported a significant correlation between protection of patients' privacy and the level of their satisfaction [12].

One of the reasons behind patient dissatisfaction with care services is nurses' poor communication with patients and their failure to educate patients about the aim(s) of hospitalization, equipment, and procedures [17]. The results of our literature review revealed that there are handful studies on the effects of different interventions on patient satisfaction. This study was conducted to examine the effects of the FCOP on satisfaction with healthcare services among patients with coronary artery disease.

## 2. Methods

This was a clinical randomized trial. Patients with coronary artery disease hospitalized in a the coronary care unit of a teaching hospital affiliated to Zanjan University of Medical Sciences, Zanjan, Iran, as well as their family members formed the study population. The inclusion criteria were having an age of 35–80, suffering from myocardial infarction or unstable angina, being completely alert and able to communicate, having a stable hemodynamic status, having no acute pain, having no previous history of hospitalization in critical care units, and having no life-threatening dysrhythmia. Patients were excluded if they died lost consciousness, developed problems which affected their communication ability, and opted for withdrawing from the study. Family members were included in the study provided that they were literate and actively involved in giving care to their patients. Family members who failed to participate in the FCOP or opted for withdrawing from the study were excluded. After conducting a pilot study, the study sample size was determined to be 80. Study participants were conveniently recruited and

randomly allocated to either the control or the experimental groups. We used the alternation technique in order to prevent the exposure of patients in the control group with educations provided to the patients in the experimental group. Accordingly, in odd weeks of month, patients were allocated to the experimental group while in the even weeks; patients were allocated to the control group.

This study was registered in the Iranian Registry of Clinical Trials with the registry code of IRCT2014030516843N2. Moreover, the study was approved by the Ethics Committee of Zanjan University of Medical Sciences, Zanjan, Iran, with the approval code of ZUMS.REC.1392.52. Before implementing the intervention, the aim of the study was explained to the participants and their informed consent was obtained. Moreover, they were assured that withdrawal from the study was voluntary and that their information would be managed confidentially.

Initially, the demographic data of patients and their family members were collected simultaneously in both study groups by using a demographic questionnaire. The content validity of the demographic questionnaire was assessed and confirmed by a panel of ten faculty members affiliated to Zanjan University of Medical Sciences, Zanjan, Iran. Then, the orientation program was implemented for patients and their family members in the experimental group according to an orientation checklist. After providing verbal educations, we provided patients and their family members with a booklet containing the same educations in the written form together with embedded pictures. We reviewed the content of the booklet with patients and family members, answered their questions, and clarified any probable ambiguities. The length of educations was 40–60 minutes. In order to ensure family members involvement in the education process, we asked them to refer to the study setting every morning and remind patients of the educations. We supervised the process of providing education to patients by their family

members in order to ensure the accuracy of educations and clarify probable ambiguities. Patients in the control group solely received routine care.

Patient satisfaction was assessed by using the Patient Satisfaction Instrument (PSI). The validity and the reliability of the PSI had been assessed and confirmed in previous studies [13 and 18]. The PSI contains 26 items in three subscales including technical professional care (seven items), trust (thirteen items), and patient education (six items). PSI items are scored on a five-point Likert scale from 1 (Completely disagree) to 5 (Completely agree). Twelve items are scored reversely. Scores of less than 78, 78–104, and greater than 104 are interpreted as dissatisfaction, moderate satisfaction, and complete satisfaction, respectively. Similarly, the scores representing dissatisfaction, moderate satisfaction, and complete satisfaction in the three subscales of technical professional care, trust, and patient education were respectively equal to ‘less than 21, 21–28, and greater than 28’, ‘less than 39, 39–52, and greater than 52’, and ‘less than 18, 18–24, and greater than 24’ [13]. Literate patients completed the questionnaires individually while for illiterate patients, questionnaires were completed by using the interview technique.

Study data were analyzed by employing the SPSS<sub>17</sub> software as well as the Kolmogorov-Smirnov, the independent-samples t, and the Fisher’s exact tests.

### 3. Results

Primarily, the Kolmogorov-Smirnov test was done for examining the distribution of the study variables. The results of this test showed that all the variables had normal distribution. Therefore, parametric statistical tests were used for data analysis.

Table 1 and 2 show the demographic characteristics of the participating patients and their family members, respectively. Most of the participating patients were male (51.2%) and aged higher than 55 years (68.75%).

After the study, all patients in the experimental group reported having either complete (17.5%) or moderate (82.5%) satisfaction and none of them were dissatisfied. However, in the control

Table 1: Patients' demographic characteristics

| Variables                           | Groups                | Control<br>N (%) | Experimental<br>N (%) | P value<br>(The Fisher's exact test) |
|-------------------------------------|-----------------------|------------------|-----------------------|--------------------------------------|
| Gender                              | Female                | 17 (42.5)        | 22 (55)               | 0.371                                |
|                                     | Male                  | 23 (57.5)        | 18 (45)               |                                      |
| Age                                 | 35-45                 | 2 (5)            | 8 (20)                | 0.117                                |
|                                     | 46-55                 | 7 (17.5)         | 8 (20)                |                                      |
|                                     | 56-65                 | 12 (30)          | 11 (27.5)             |                                      |
|                                     | 66-75                 | 9 (22.5)         | 10 (25)               |                                      |
|                                     | 76-80                 | 10 (25)          | 3 (7.5)               |                                      |
| Marital status                      | Single                | 0 (0)            | 1 (2.5)               | 0.781                                |
|                                     | Married               | 31 (77.5)        | 31 (77.5)             |                                      |
|                                     | Divorced              | 0 (0)            | 1 (2.5)               |                                      |
|                                     | Widowed               | 9 (22.5)         | 7 (17.5)              |                                      |
| Employment                          | White-collar worker   | 6 (15)           | 6 (15)                | 0.393                                |
|                                     | Blue-collar worker    | 7 (17.5)         | 9 (22.5)              |                                      |
|                                     | Other                 | 5 (12.5)         | 3 (7.5)               |                                      |
|                                     | Housewife             | 15 (37.5)        | 20 (50)               |                                      |
|                                     | Unemployed            | 7 (17.5)         | 2 (5)                 |                                      |
| Education                           | Illiterate            | 20 (50)          | 15 (37.5)             | 0.431                                |
|                                     | Below diploma         | 10 (25)          | 15 (37.5)             |                                      |
|                                     | Diploma or higher     | 10 (25)          | 10 (25)               |                                      |
| Insurance                           | Have                  | 40 (100)         | 39 (97.5)             | 1.00                                 |
|                                     | Do not have           | 0 (0)            | 1 (2.5)               |                                      |
| Supplementary insurance             | Have                  | 5 (12.5)         | 7 (17.5)              | 0.755                                |
|                                     | Do not have           | 35 (87.5)        | 33 (82.5)             |                                      |
| History of previous hospitalization | Have                  | 22 (55)          | 18 (45)               | 0.503                                |
|                                     | Do not have           | 18 (45)          | 22 (55)               |                                      |
| Diagnosis                           | Angina pectoris       | 23 (57.5)        | 23 (57.5)             | 1.00                                 |
|                                     | Myocardial infarction | 17 (42.5)        | 17 (42.5)             |                                      |

Table 2: Demographic characteristics of family members in the experimental group

| Variables           | Groups              | N (%)     |
|---------------------|---------------------|-----------|
| Gender              | Female              | 21 (52.5) |
|                     | Male                | 19 (47.5) |
| Marital status      | Single              | 20 (50)   |
|                     | Married             | 20 (50)   |
| Employment          | White-collar worker | 10 (25)   |
|                     | Blue-collar worker  | 4 (10)    |
|                     | Other               | 3 (7.5)   |
|                     | Housewife           | 15 (37.5) |
|                     | Unemployed          | 8 (20)    |
| Education           | Below diploma       | 5 (12.5)  |
|                     | Diploma or higher   | 35 (87.5) |
| Kinship             | Spouse              | 6 (15)    |
|                     | Children            | 24 (60)   |
|                     | Parents             | 1 (2.5)   |
|                     | Sibling             | 2 (5)     |
|                     | Grandchildren       | 3 (7.5)   |
|                     | Daughter-in-law     | 4 (10)    |
| Living with patient | Yes                 | 27 (67.5) |
|                     | No                  | 13 (32.5) |

group, 35 patients (87.5%) were moderately satisfied, five patients (12.5%) were dissatisfied, and none of them were completely satisfied with care services.

The results of the independent-samples t test revealed that after implementing the study intervention, the mean of PSI score in the experimental group was significantly higher than that of the control group (97.88 vs. 85.15;  $p < 0.0001$ ). Moreover, the scores of the three sub-scales of technical professional care, trust, and patient education in the experimental group were significantly higher than the control group ( $p < 0.05$ ; Table 3).

#### 4. Discussion

The findings of the current study revealed that implementing the FCOP enhanced satisfaction with care services among patients with

coronary artery disease. We also found that the FCOP enhanced patient satisfaction in the three PSI sub-scales of technical professional care, trust, and patient education. Overall, our participating patients had moderate satisfaction with care services. However, Lee et al. (2008) reported that the rate of satisfaction with primary care services among patients with myocardial infarction in Canada was 91.7% [19].

Our participants' greatest satisfaction was with technical professional care. This is in line with

On the other hand, the lowest level of satisfaction was related to the patient education sub-scale. This is in congruence with the findings of previous studies [13, 18, and 20]. Patient education is among nurses' main professional responsibilities and patients tend to receive educations from their nurses [21]. However, our patients had low satisfaction with patient education, denoting nurses' poor patient education practice. This finding can be attributed to the fact that the participating patients evaluated all nurses who had provided

Table 3: Patient satisfaction with the three sub-scales of technical professional care, trust, and patient education

| Subscales                   | Groups       | Complete satisfaction |      | Moderate satisfaction |      | dissatisfaction |      | P value (the independent-samples t test) |
|-----------------------------|--------------|-----------------------|------|-----------------------|------|-----------------|------|--|
|                             |              | N                     | %    | N                     | %    | N               | %    |  |
| Technical professional care | Control      | 3                     | 7.5  | 27                    | 67.5 | 10              | 25   | 0.01                                     |
|                             | Experimental | 12                    | 30   | 23                    | 57.5 | 5               | 12.5 |  |
|                             | Total        | 15                    | 18.8 | 50                    | 62.5 | 15              | 18.8 |  |
| Trust                       | Control      | 2                     | 5    | 28                    | 70   | 10              | 25   | < 0.0001                                 |
|                             | Experimental | 12                    | 30   | 26                    | 65   | 2               | 5    |  |
|                             | Total        | 14                    | 17.5 | 54                    | 67.5 | 12              | 15   |  |
| Patient education           | Control      | 2                     | 5    | 27                    | 67.5 | 11              | 27.5 | 0.008                                    |
|                             | Experimental | 8                     | 20   | 28                    | 70   | 4               | 10   |  |
|                             | Total        | 10                    | 12.5 | 55                    | 68.8 | 15              | 18.8 |  |

the findings reported by Hajinezhad et al. (2007) and Wolf et al. (2003) [18 and 20]. However, Joulai et al. (2011) reported that patients' greatest satisfaction was related to the trust sub-scale of the PSI [13]. Our participants' great satisfaction with technical professional care can be attributed to the overriding importance of technical care behaviors from the perspective of nurses which makes them focus on patient care despite having heavy workload. Moreover, compared with other aspects of patient satisfaction, technical professional care is more tangible and observable and hence, it can be directly assessed and easily scored by patients. Arefi and Talaei (2010) also found that patients' greatest satisfaction was related to nurses' timely attendance at their bedside [14], confirming nurses' greater attention to technical care behaviors.

care to them during the course of their hospitalization.

## 5. Conclusions

Study findings indicate that the FCOP can enhance patient satisfaction. Given the low score of the patient education subscale of patient satisfaction, greater attention should be paid by nurses to patient education.

One of the study limitations was that the study instrument mainly assessed patient satisfaction with nursing care. Future studies are recommended to use more comprehensive instruments for assessing different aspects of patient satisfaction. Moreover, satisfaction is an abstract concept which can be affected by patients' mood and psychological state. Accordingly, study findings might have been

affected by patients' mood and psychological state.

## 6. Acknowledgments

This article was part of a Master's thesis which had been funded by the Research Council of Zanjan University of Medical Sciences, Zanjan, Iran. We are genuinely grateful to all the nurses, patients, and family members who helped us conduct this study.

## References

1. Moser D.K, Chung M, McKinley S, Riegel B, An K, Cherrington C.C, et al. Critical care nursing practice regarding patient anxiety assessment and management. *Intensive Crit Care Nurs.*2003; 19(5):276-288.
2. Chaboyer W, Thalib L, Alcorn K, Foster M. The effect of an ICU liaison nurse on patients and family's anxiety prior to transfer to the ward: An intervention study. *Intensive Crit Care Nurs.*2007;23 (6): 362-369.
3. Chien W.T, Chiu Y.L, Lam L.W, Ip W.Y. Effects of a needs based education programme for family careers with a relative in an intensive care unit: A quasi-experimental study. *Int J Nurs Stud.*2006; 43(1): 39-50.
4. Rabie Siahkali S, Pourmemari MH, Khaleghdoost Mohamadi T, Eskandari F, Avazeh A. study on effective factors on patients' family members anxiety in intensive care units. *Journal of zanjan university of medical sciences.*2010; 18 (70): 91-101.
5. Kalda R, Oona M, Maaroos H, Lember M. Patient evaluation on family doctors' family orientation. *Patient Educ Couns.*2005; 56(3):296-301.
6. Bailey J.J, Sabbagh M, Loiselle C.G, Boileau J, McVey L. Supporting families in the ICU: A descriptive correlational study of informational support, anxiety, and satisfaction with care. *Intensive Crit Care Nurs.*2010; 26(2): 114-122.
7. Ishikawa H, Takayama T, Yamazaki Y, Seki Y, Katsumata N. Physician-patient communication and patient satisfaction in Japanese cancer consultations. *Social Science & Medicine.* 2002;55(2):301-311.
8. Shaw W.S, Jane Woiszwillo M, Krupat E. Further validation of the Patient-Practitioner Orientation Scale (PPOS) from recorded visits for back pain. *Patient education and counseling.* 2012;89(2):288-291.
9. Mallinger J.B, Griggs J.J, Shields C.G. Patient-centered care and breast cancer survivors' satisfaction with information. *Patient education and counseling.* 2005;57(3):342-349.
10. Lubart E, Leibovitz A, Shapira A, Tischenko O, Peled Z, Baumohl Y, et al. Satisfaction with the care of institutionalized psychogeriatric patients, as reflected by a survey of their relatives. *Archives of gerontology and geriatrics.* 2004;38(1):45-50.
11. Farahani M, Kashaninia Z, Hoseini M, Biglarian A. The effect of education of communication skills to nurses on patients satisfaction. *Nursing research.* 2007;1(3):47-54.
12. Dehghannayyeri N, Aghajani M. Relationship between privacy and patients satisfaction in emergency ward. *Hayat.*16(1):13-22.
13. Joulai S, Hajibabai F, Jafarjalal E, Bahrani N. Evaluation of patients satisfaction from nursing care. *Hayat.* 2011;17(1):35-44.
14. Arefi M, Talai N. Evaluation of patients. *Payavarde salamat.* 2010;4(1,2):97-103.
15. Mohammadi A, Vanaki Z, Mohammadi A. Effect of implementation of motivational program based on "expectancy theory" by head nurses on patients' satisfaction. *Hayat.* 2012;18(4):47-60.
16. Chan R.J, Webster J, Marquart L. A systematic review: The effects of orientation programs for cancer patients and their family/carers. *International Journal of nursing studies.* 2012;49(12):1558-1567.
17. Stricker K.H, Niemann S, Bugnon S, Wurz J, Rohrer O, Rothen H.U. Family satisfaction in the intensive care unit : cross-cultural adaptation of a questionnaire. *Journal of critical care.* 2007;22(3):204-211.
18. Hajinezhad M, Rafii F, Jafarjalal E, Haghani H. Relationship between nurse caring behaviors from patients' perspectives & their satisfaction. *Iran Journal of nursing.* 2007;20(49):73-83.
19. Lee D.S, Tu J.V, Chong A, Alter D.A. Patient satisfaction and its relationship with quality and outcomes of care after acute myocardial infarction. *Circulation.* 2008;118(19):1938-1945.
20. Wolf Z.R, Miller P.A, Devine M. Relationship between nurse caring & patient satisfaction in patient undergoing invasive cardiac procedures. *Med Surg Nurs.* 2003;12(6):391-397.
21. Asilioglu K, Celik S.S. The effect of preoperative education on anxiety of open cardiac surgery patients. *Patient education and counseling.* 2004;53(1):65-70.
22. Kristjanson L.J. Validity and reliability testing of the Famcare scale: measuring family satisfaction with advanced cancer care. *Social science & medicine.* 1993;36(5):693-701.